

Figure 1. Plan of Niah Cave showing the location of the main entrances (adapted from Majid 1982).

39,600±1000 and 41,500±1000 BP respectively. If these dates are indicative of the age of the deep skull, then Niah provides the earliest evidence of modern humans in Southeast Asia.

EVIDENCE FOR ECOLOGICAL CHANGE

The dense accumulation of mammalian bones at Niah appears to be primarily due to human subsistence activities. There is ample evidence of bone charring in the excavated remains, but indications of bone modification and butchering marks are quite rare. The fauna is extensive, with 59 species of mammals being represented, although this includes 13 species of bats and other species of micromammals, many of which are unlikely to be present as the result of human hunting activities (Medway 1966a, 1977).

Harrison (1961) suggested that human hunting during the late Pleistocene had a profound effect on the fauna of Borneo. He used the following evidence from Niah to support his argument:

1. At least one species of mammal found at Niah, the giant pangolin (*Manis palaeojavanica*) has become entirely extinct (Hooijer 1960a; Harrison *et al.* 1961) (Table 1)
2. Several species of large mammals found at Niah and present in other parts of southeast Asia today are now extinct in Borneo (i.e., *Panthera tigris* and *Tapirus indicus*¹) (Medway 1959a, 1960, 1964a, 1977) (Table 1)
3. Some species found at Niah are rare or absent in the local area today (i.e., *Dicerorhinus sumatrensis*, *Elephas maximus*, and *Pongo pygmaeus*)

Table 1. List of large mammal species (excluding domesticates) from Niah^a

Species represented at Niah		Status in Borneo today
Primates		
<i>Pongo pygmaeus</i>	Orang-utan	Present
<i>Hylobates muelleri</i>	Gibbon	Present
<i>Trachypithecus cristatus</i>	Silvered leaf monkey	Present
<i>Presbytis</i> spp. (? <i>P. rubicunda</i>)	Leaf monkey	Present
<i>Macaca fascicularis</i>	Long-tailed macaque	Present
<i>Macaca nemestrina</i>	Pig-tailed macaque	Present
<i>Nycticebus coucang</i>	Slow loris	Present
Pholidota		
<i>Manis palaeojavanica</i>	Giant pangolin	Extinct
<i>Manis javanica</i>	Malay pangolin	Present
Carnivora		
<i>Ursus malayanus</i>	Sun bear	Present
<i>Panthera tigris</i>	Tiger	Absent
<i>Pardofelis nebulosa</i>	Clouded leopard	Present
<i>Mustela nudipes</i>	Malayan weasel	Present
<i>Melogale everetti</i>	Bornean ferret-badger	Mt. Kinabalu only
<i>Lutra sumatrana</i>	Hairy-nosed otter	Present
<i>Viverra zangalunga</i>	Malay civet	Present
<i>Arctictis binturong</i>	Bearcat	Present
<i>Hemigalus derbyanus</i>	Banded palm civet	Present
<i>Herpestes</i> sp.	Mongoose	Present
Perissodactyla		
<i>Tapirus indicus</i>	Malayan tapir	Absent
<i>Dicerorhinus sumatrensis</i>	Sumatran rhinoceros	Present, but rare
Artiodactyla		
<i>Sus barbatus</i>	Bearded pig	Present
<i>Tragulus napu</i>	Large Malay mouse-deer	Present
<i>Tragulus javanicus</i>	Lesser Malay mouse-deer	Present
<i>Muntiacus muntjak</i>	Barking deer	Present
<i>Cervus unicolor</i>	Sambar	Present
<i>Bos javanicus</i>	Banteng	Present
Proboscidea		
<i>Elephas maximus</i>	Asian elephant	Present in Sabah only

^a Sources: Medway (1959a, 1959b, 1960, 1964a, 1966b, 1977); Harrison (unpublished); Corbet & Hill (1992).

4. Several forms are larger in size, especially at deeper levels in the cave deposits.

By contrast, Medway (1977; Cranbrook 1988) has argued that the impact of hunting on the fauna was insignificant, and that variations in the mammalian community structure during the late Quaternary were due to ecological changes. Increasing evidence from a number

of inter-related lines of research tend to support this latter view:

1. The most common mammal remains from Niah (presumably the most commonly hunted mammals) are the orang-utan, several species of monkeys, and the bearded pig, which are still frequent in Borneo at present, while the large mammals that are now rare or

Table 2. Frequency of large mammal species at Niah

Very common > 200 specimens	Quite common >20 specimens		Rare < 20 specimens
<i>Pongo pygmaeus</i>	<i>Trachypithecus cristatus</i>	<i>Hylobates muelleri</i>	<i>Viverra tangalunga</i>
<i>Presbytis</i> spp.	<i>Bos javanicus</i>	<i>Macaca nemestrina</i>	<i>Arctictis binturong</i>
<i>Macaca fascicularis</i>		<i>Manis palaeojavanica</i>	<i>Hemigalus derbyanus</i>
<i>Sus barbatus</i>		<i>Manis javanica</i>	<i>Herpestes</i> sp.
		<i>Ursus malayanus</i>	<i>Muntiacus muntjak</i>
		<i>Panthera tigris</i>	<i>Cervus unicolor</i>
		<i>Pardofelis nebulosa</i>	<i>Tragulus napu</i>
		<i>Mustela nudipes</i>	<i>Tragulus javanicus</i>
		<i>Melogale everetti</i>	<i>Tapirus indicus</i>
		<i>Lutra sumatrana</i>	<i>Dicerorhinus sumatrensis</i>
			<i>Elephas maximus</i>

absent locally are also very scarce in the collections (Table 2). This is hardly the pattern to be expected if humans on Borneo were responsible for exterminating species through over-hunting.

- The impact that human hunting has on local faunas is related to the extent to which the fauna has co-evolved with hominid predation in the past. Where humans have come into contact for the first time with highly endemic faunas, such as in the New World, Australia or Madagascar, hunting has led to major extinctions. In contrast, where communities have co-evolved with hominids for a long time, as in Eurasia and Africa during the Pleistocene, levels of extinction are much lower. The mammal fauna on Borneo in the late Pleistocene was typically Indo-Malayan, apparently with few endemic species, and one with a long evolutionary history of association with hominids. In addition, archaeological and historical evidence suggests that human populations on Borneo may have been quite sparse (Bellwood 1990, 1993), in which case hunting may have had little impact on the fauna.
- Evidence from other parts of Southeast Asia suggest that eustatic fluctuations had a pronounced effect on the ecology and climate of northern Borneo during the late Pleistocene. The mean annual temperature on Borneo at the time of the last glacial maximum, around 18,000 BP, might have been 5-6°C lower than present (20-22°C), and precipitation was reduced and more seasonally distributed (Peterson 1969; Haile 1971; Biswas 1973; Verstappen 1975; Medway 1977). With a sea level probably as much as 120 m below that at present, Niah would have been more

than 200 km inland from its present position, which is now only 16 km from the coast. The drier, more seasonal, forest would have provided a more suitable biome for large herbivorous mammals, along with their dependent predators (Medway 1977). With the establishment of the present climatic regime during the Holocene, and the development of ever-wet tropical rainforest, suitable habitats for these large mammals diminished, leading to declining numbers and extinction.

- The Niah Cave fauna itself indicates ecological changes. Larger dental or inferred body size of the subfossil specimens relative to extant conspecifics has been reported in the orang-utan, macaques, leaf monkeys, murids, Sumatran rhinoceros and barking deer (Hooijer 1961, 1962; Medway 1959b, 1964b, 1966a, 1966b; Cranbrook 1986). This is presumably associated with a purportedly cooler early Quaternary climate. The presence of two exclusively montane forest species, the lesser gymnure (*Hylomys suillus*) and ferret badger (*Melogale everetti*), also supports this view (Medway 1964a, 1977).
- Heaney (1986) has shown that the rate of extinction of mammals by attrition on the islands of southeast Asia following inundation of the Sunda shelf to be correlated with island size. Such extinctions would have had a minor impact on an island the size of Borneo, but may have been sufficient to account for the low level of impoverishment indicated by the archaeological record at Niah and other prehistoric cave sites in Borneo.

Table 3. The percentage of primates (cranio-dental specimens) by level from the West Mouth at Niah

Depth	N	<i>Presbytis</i> spp.	<i>Macaca</i> <i>fascicularis</i>	<i>Macaca</i> <i>nemestrina</i>	<i>Trachypith.</i> <i>cristatus</i>	<i>Pongo</i> <i>pygmaeus</i>	<i>Hylobates</i> <i>muelleri</i>
Unknown	27	18.5	18.5	7.4	18.5	37.0	0.0
0-12"	23	26.1	8.7	4.3	21.7	39.1	0.0
12-24"	37	24.3	13.5	2.7	8.1	51.4	0.0
24-36"	50	18.0	24.0	4.0	28.0	24.0	2.0
36-48"	28	35.7	10.7	0.0	0.0	53.6	0.0
48-60"	17	11.8	23.5	0.0	11.8	52.9	0.0
+ 60"	35	34.3	11.4	2.9	0.0	51.4	0.0
Total	100.0%	24.4%	16.1%	3.2%	13.4%	42.4%	0.5%
N	217	53	35	7	29	92	1

Table 4. The density of extant primates in Borneo compared with the frequency of subfossil primates from West Mouth, Niah

	Kutai Nature Reserve, Borneo ^a		Niah, West Mouth
	Individuals per km ²	Frequency	Frequency
<i>Pongo pygmaeus</i>	4.0	7.9%	42.4%
<i>Hylobates muelleri</i>	14.6	29.0%	0.5%
Leaf monkeys	20.4	40.6%	37.8%
<i>Macaca nemestrina</i>	5.8	11.5%	3.2%
<i>Macaca fascicularis</i>	5.5	10.9%	16.1%

^a Data from Rodman (1978), Waser (1987).

PRIMATE MATERIAL FROM NIAH CAVE

My preliminary research on the subfossil primates from Niah contributes new information that is pertinent to understanding the nature of the ecological changes that took place in Borneo during the late Quaternary. Along with pigs and squirrels, primates are the most common mammals at Niah. Eight species of primates are represented, including all genera currently living on Borneo, with the exception of *Tarsius* and *Nasalis*.

The most common primate at Niah West Mouth is *Pongo pygmaeus*, which comprises 42.4% of the total from the site. This is followed in importance by *Presbytis* spp. (24.4%), *Macaca fascicularis* (16.1%) and *Trachypithecus cristatus* (13.4%). *Macaca nemestrina* and *Hylobates muelleri* are, by contrast, quite rare. The distribution of these species with increasing depth in the cave does not appear to change significantly, but it should be noted that the sample sizes are relatively small (Table 3). Medway (1959b) has suggested that there was an abrupt increase in the relative frequency of arboreal

animals, such as monkeys and squirrels, above the 48 inch level, which he associates with the advent of a more sophisticated technology, including greater use of bone as a raw material, especially in the manufacture of projectile tips.

However, raw data on the absolute numbers of primate specimens provides no real clues to how intensively these primates were being exploited for food. Density information for living primates gives us some indication of the relative abundance of different primates in a modern setting (Table 5). If these frequencies are representative of primate communities on Borneo in the past, then we can gain some idea of hunting preference from them. The data seem to suggest that orang-utans and leaf monkeys are over-represented in the collections from Niah, while the macaques, and especially the gibbons, are dramatically under-represented. This different representation may reflect either food preferences by prehistoric human populations or more likely that orang-utans and leaf monkeys are easier prey to catch.

Table 5. Percentage of primates (cranio-dental specimens) from the major sites at Niah.

Site	N	<i>Presbytis</i> spp.	<i>Macaca</i> <i>fascicularis</i>	<i>Macaca</i> <i>nemestrina</i>	<i>Trachypith.</i> <i>cristatus</i>	<i>Pongo</i> <i>pygmaeus</i>	<i>Hylobates</i> <i>muelleri</i>
West Mouth	217	24.4	16.1	3.2	13.4	42.4	0.5
Lobang Angus	114	20.2	50.9	0.9	14.9	12.3	0.9
Gan Kira	38	26.3	28.9	2.6	7.9	31.6	2.6

Table 6. Comparison of occlusal area (mesiodistal length x buccolingual breadth) of upper and lower cheek teeth combined in modern and subfossil orang-utans from Borneo and Sumatra^a

Standard deviation intervals from the mean values for extant orang-utans

	N	-3	-2	-1	+1	+2	+3	+4	+5
Extant	822	4	136	304	235	113	29	1	0
<i>Pongo</i>	100%	0.5%	16.5%	37.0%	28.6%	13.7%	3.5%	0.1%	0%
Padang	2672	0	72	592	917	655	323	87	26
<i>Pongo</i> ^b	100%	0%	2.7%	22.2%	34.3%	24.5%	12.1%	3.3%	1.0%
Niah Cave	65	0	3	24	30	7	1	0	0
<i>Pongo</i>	100%	0%	4.6%	36.9%	46.2%	10.8%	1.5%	0%	0%

^a Data obtained from Hooijer (1948) and Harrison (unpublished).

^b The subfossil orang-utans from late Quaternary sites in the Padang Highlands of Sumatra have been described by Hooijer (1948). The results presented here confirm Hooijer's contention that the dental remains from the Padang Highlands are significantly larger than the teeth of extant orang-utan.

Aside from this, the distribution of different primate species in the collections from the various cave mouths at Niah differ significantly from each other: orang-utans represent 42.4% of all primates at the West Mouth, but only 12.3% at Lobang Angus, while the long-tailed macaque (*Macaca fascicularis*) represents 16.1% and 50.9% respectively (Table 6). A simple explanation is not possible. The composition of the forest around Lobang Angus, which opens onto the eastern side of Subis, may have been somewhat distinct from that surrounding the west facing Main Mouth, thereby allowing human hunters access to subtly different mammalian communities. However, this suggestion needs to be tested further with comparisons of the remaining fauna.

The suggestion that several species of primates have undergone size reduction in Borneo during the Quaternary, presumably as a response to increased ambient temperatures (Hooijer 1961, 1962), is an important piece of evidence used in support of changed ecological conditions. The orang-utans and the long-tailed macaques provide the best samples from Niah to test this inference. Orang-utans have a quite limited distribution in Sarawak

today and are not found in the vicinity of Niah (Figure 2). This is probably as a result of recent hunting pressure and disturbance (Medway 1977). When the dental size of extant orang-utans is compared with those from Niah, the latter are not significantly different. A re-examination of the dental size of orang-utans from Niah show that the teeth tend to be slightly larger on average than those of modern animals, but all specimens fall well within the range for extant orang-utans (Tables 6 and 7). This finding contradicts the conclusion of Hooijer (1961) that the subfossil orang-utans from Borneo were relatively larger than their modern counterparts.

The long-tailed macaques provide a more convincing case for significant size reduction: the cheek teeth of the subfossil forms are on average 13% larger (based on occlusal area) than those of their modern conspecifics from Borneo (Table 8). In addition, there appears to have been a gradual reduction in overall size of the dentition through time (Figure 3). A recent study of skull length (used as a measure of overall size) in *Macaca fascicularis* has shown a gradual increase with distance from the Equator (Fooden and Albrecht 1993). They explain this




Figure 2. The recent distribution of orang-utans in Sarawak in relation to Niah Cave (adapted from Reynolds 1967).

as a consequence of Bergmann's rule, in which increased body size occurs as an adaptive response to cooler climates. It should be noted that the subfossil macaques from Niah correspond in dental size with those that inhabit regions of Thailand today which have mean annual temperatures similar to that inferred for northern Borneo during the last glacial maximum.

CONCLUSIONS

There is growing evidence to support the hypothesis that ecological change, associated with global eustatic fluctuations, had a more profound impact on the diversity and distribution of the late Quaternary large mammal fauna of Borneo than did environmental disturbance by humans. A re-examination of the primate fauna from

Niah Cave contributes new evidence pertinent to understanding ecological changes that occurred in Borneo during the last 40,000 years. The major findings of this study can be summarized as follows:

1. The orang-utan, leaf monkeys and macaques represent significant components of the modern fauna of Borneo, despite what appears from the archaeological evidence at Niah to have been a pattern of preferential hunting of these species by humans throughout the late Quaternary;
2. *Pongo pygmaeus* and *Presbytis* spp. appear to be over-represented in the collections at Niah, at least when compared with modern patterns of abundance, and this may reflect either food preferences or limitations in the hunting capabilities of prehistoric humans;

Table 7. Comparison of the occlusal area (mesiodistal length x buccolingual breadth) of the permanent teeth of the orang utan sample from West Mouth, Niah, compared with the standard deviation intervals from the means of the corresponding teeth of extant orang-utans, arranged by their depth in the cave.

	Depth at Niah	N	Minimum S.D. interval	Maximum S.D. interval	Mean S.D. interval
Extant orang-utans		822	-2.50	+3.50	0.0
Subfossil orang-utans	0"-12"	5	-0.87	+0.34	-0.12
	12"-24"	13	-0.96	+0.95	-0.27
	24"-36"	8	-0.96	+1.42	+0.15
	36"-48"	12	-0.34	+1.62	+0.65
	48"-60"	2	+0.47	+0.77	+0.62
	60"-72"	7	-0.26	+2.00	+0.79
	72"-84"	1	-0.27	-0.27	-0.27
	Below 84"	2	+0.84	+1.03	+0.94
All levels		57	-1.29	+2.00	+0.19

Data from Hooijer (1948) and Harrison (unpublished)

Table 8. Relative difference in occlusal areas (mesiodistal length x buccolingual breadth) of cheek teeth in extant *Macaca fascicularis* from Borneo and subfossil *M. fascicularis* from Niah.

	Mean area in extant form	Mean area in subfossils	% Difference
Upper P4	23.8	26.7	+12.2%
Upper M1	35.7	38.8	+8.7%
Upper M2	44.5	50.5	+13.5%
Upper M3	41.6	47.2	13.5%
Lower P4	19.9	23.6	+18.6%
Lower M1	27.8	32.0	+15.1%
Lower M2	39.5	44.2	+11.9%
Lower M3	45.7	51.1	+11.8%
Mean Difference			+13.2%

- Differences in the proportions of primates from the various cave mouths at Niah may reflect subtle ecological variations in the forest surrounding the Subis limestone massif, thereby allowing human hunters access to different resources;
- The subfossil orang-utans from Niah do not differ significantly in dental size from their extant counterpart, contrary to Hooijer (1960b);
- The long-tailed macaques (*Macaca fascicularis*) from Niah are significantly larger in dental size than their modern conspecifics from Borneo, but they apparently decrease in size through time; and
- The subfossil macaques from Niah have teeth similar in size to those of *Macaca fascicularis* populations

living today in Thailand where the mean annual temperatures correspond to those inferred for northern Borneo during the last glacial maximum, at about 18,000 BP.

ACKNOWLEDGMENTS

I would like to thank the following individuals and institutions for allowing me access to materials in their care: Peter Kedit and Ipoi Datan, Sarawak Museum, Kuching; Jon de Vos, Rijksmuseum van Natuurlijke Historie, Leiden; Jerry Hooker and Peter Andrews, The Natural History Museum, London; Guy Musser and Wolfgang Fuchs, American Museum of Natural History, New York.

DENTAL SIZE OF *M. FASCICULARIS* FROM NIAH




Figure 3. The dental size of subfossil *Macaca fascicularis* from Niah compared with that of extant conspecifics from Borneo. The samples are arranged according to their depth in the cave; sample sizes are given in parentheses. The x-axis represents the standard deviation intervals from the mean values of the occlusal areas (mesiodistal length x buccolingual breadth) of all upper and lower cheek teeth in extant *M. fascicularis*. Solid bars are ranges; diamonds are mean values. Note that the cheek teeth of subfossil macaques from Niah are larger at deeper levels in the cave deposits, and that they gradually decrease through time to the present.

Bernard Maloney deserves special thanks for greatly improving the clarity of the text and for preparing the figures. I am also grateful to Peter Andrews, Ipoi Datan, John Krigbaum, Michael Rampino, Chris Stringer and Jon de Vos for discussions and helpful suggestions. This research was supported by a travel grant from New York University.

NOTE

1. Evidence from archaeological sites in Sabah demonstrate that the Javan rhinoceros (*Rhinoceros sondaicus*) and the dhole (*Cuon alpinus*) have also become extinct on Borneo since the late Quaternary (Cranbrook 1986, 1988).

REFERENCES

Bellwood, P. 1990. From late Pleistocene to early Holocene in Sundaland. In C. Gamble and O. Soffer (eds.), *The World*

- at 18,000 BP, Vol. 2: *Low Latitudes*, pp. 255-263. London: Unwin Hyman.
- 1993. Cultural and biological differentiation in Peninsular Malaysia: the last 10,000 years. *Asian Perspectives* 32:37-60.
- Biswas, B. 1973. Quaternary changes in sea level in the South China Sea. *Bulletin of the Geological Society of Malaysia* 6:229-256.
- Brooks, S.T.; Hegler, R. and Brooks, R.H. 1977. Radiocarbon dating and palaeoserology of a selected burial series from the Great Cave of Niah, Sarawak, Malaysia. *Asian Perspectives* 20:21-31.
- Brothwell, D.R. 1960. Upper Pleistocene human skull from Niah Caves, Sarawak. *Sarawak Museum Journal* 9:323-350.
- Chesner, C.A.; Rose, W.I.; Deino, A.; Drake, R. and Westgate, J.A. 1991. Eruptive history of Earth's largest Quaternary caldera (Toba, Indonesia) clarified. *Geology* 19:200-203.
- Corbet, G.B. and Hill, J.E. 1992. *The Mammals of the Indomalayan Region: A systematic Review*. Oxford: Oxford University Press.
- Cranbrook, Earl of 1986. A review of fossil and prehistoric remains of rhinoceroses in Borneo. *Sabah Museum Journal* 1:50-110.
- 1988. The contribution of archaeology to the zoogeography of Borneo, with the first record of a wild canid of early Holocene age. *Fieldiana Zoology* 42:1-7.
- Fooden, J and Albrecht, G.H. 1993. Latitudinal and insular variation of skull size in crab-eating macaques (Primates, Cercopithecidae: *Macaca fascicularis*). *American Journal of Physical Anthropology* 92:521-538.
- Haile, N.S. 1971. Quaternary shorelines in West Malaysia and adjacent parts of the Sunda shelf. *Quaternaria* 15:333-343.
- Harrisson, B. 1967. A classification of stone age burials from Niah great cave, Sarawak. *Sarawak Museum Journal* 15:126-200.
- Harrisson, T. 1957. The Great Cave of Niah: a preliminary report on Bornean prehistory. *Man* 57:161-166.
- 1958. The caves of Niah: A history of prehistory. *Sarawak Museum Journal* 8:549-595.
- 1959a. Radio Carbon - C-14 datings B.C. from Niah: a note. *Sarawak Museum Journal* 9: 36-138.
- 1959b. New archaeological and ethnological results from Niah Cave, Sarawak. *Man* 59: 1-8.
- 1961. Prehistoric fauna changes and losses in Borneo. *The Malayan Nature Journal* (21st. Anniversary, Special Issue): 89-92.
- 1970. The prehistory of Borneo. *Asian Perspectives* 13: 17-45.
- 1975. Early dates for "seated" burial and burial matting at Niah Caves, Sarawak (Borneo). *Asian Perspectives* 18:161-165.
- Harrisson, T.; Hooijer, D.A. and Medway, Lord 1961. An extinct giant pangolin and associated mammals from Niah Cave, Sarawak. *Nature* 189:166.
- Heaney, L.R. 1986. Biogeography of mammals in SE Asia: estimates of rates of colonization, extinction and speciation. *Biological Journal of the Linnean Society* 28:127-165.
- Hooijer, D.A. 1948. Prehistoric teeth of man and of the orang-utan from central Sumatra, with notes on the fossil orang-utan from Java and southern China. *Zoologische Mededeelingen* 29:175-301.
- 1960a. The giant extinct pangolin (*Manis palaeojavanica* Dubois) from Niah. *Sarawak Museum Journal* 9:350-355.
- 1960b. Quaternary gibbons from the Malay Archipelago. *Zoologische Verhandelingen Museum Leiden* 46:1-42.
- 1961. The orang-utan in Niah Cave pre-history. *Sarawak Museum Journal* 9:408-421.
- 1962. Prehistoric bone: the gibbons and monkeys of Niah Great Cave. *Sarawak Museum Journal* 10: 428-449.
- Kennedy, K. A. R. 1977. The deep skull of Niah: An assessment of twenty years of speculation concerning its evolutionary significance. *Asian Perspectives* 20:32-50.
- Majid, Z. 1982. The West Mouth, Niah, in the prehistory of Southeast Asia. *Sarawak Museum Journal* 31:1-200.
- Medway, Lord 1959a. The tapir at Niah. *Sarawak Museum Journal* 9:146.
- 1959b. Niah animal bone: II (1954-8). *Sarawak Museum Journal* 9:151-163.
- 1960. The Malay Tapir in late Quaternary Borneo. *Sarawak Museum Journal* 9:356-360.
- 1964a. Post-Pleistocene changes in the Mammalian fauna of Borneo. *Studies in Speleology* 1:33-37.
- 1964b. Niah Cave bone - VII: size changes in the teeth of two rats, *Rattus sabanus* Thomas and *R. muelleri* Jentink. *Sarawak Museum Journal* 11:616-623.
- 1966a. Animal remains from Lobang Angus, Niah. *Sarawak Museum Journal* 14:185-216.
- 1966b. Niah Cave animal bone. VIII - Rhinoceros in late Quaternary Borneo. *Sarawak Museum Journal* 12:77-82.
- 1977. The Niah excavations and an assessment of the impact of early man on mammals in Borneo. *Asian Perspectives* 20:51-69.
- Nishimura, S. and Stauffer, P.H. 1981. Fission-track dating of zircons from the Serdang volcanic ash, Peninsular Malaysia. *Warta Geologi* 7:39-41.
- Petersen, R.M. 1969. Würm II climate at Niah Cave. *Sarawak Museum Journal* 17: 67-79.
- Reynolds, V. 1967. *The Apes*. New York, Harper & Row.

- Shutler, R. 1977. Tom Harrison's contribution through radiocarbon dating to the understanding of the prehistory of Southeast Asia. *Asian Perspectives* 20:9-12.
- Solheim, W.G. 1958. The present status of the 'Palaeolithic' in Borneo. *Asian Perspectives* 2:83-90.
- 1974. Borneo archaeology, past and future. *Sabah Society Journal* 6:27-37.
- 1983. Archaeological research in Sarawak, past and future. *Sarawak Museum Journal* 32:35-58.
- Spriggs, M. 1989. The dating of the island Southeast Asian Neolithic: an attempt at chronometric hygiene and linguistic correlation. *Antiquity* 63:587-613.
- Stauffer, P.H.; Nishimura, S. and Batchelor, B.C. 1980. Volcanic ash in Malaya from a catastrophic eruption of Toba, Sumatra, 30,000 years ago. In S. Nishimura (ed.), *Physical Geology of Indonesian Island Arcs*, pp.156-164. Kyoto: Kyoto University Press.
- Verstappen, H. T. 1975. On palaeoclimates and landform development in Malesia. *Modern Quaternary Research in S.E. Asia* 1:3-36.
- Waser, P. 1987. Interactions among primate species. In B.B. Smuts; D.L. Cheney; R.M. Seyfarth; R.W. Wrangham and T.T. Struhsaker (eds), *Primate Societies*, pp. 210-226. Chicago: University of Chicago Press.